

Elastic theory of quantum Hall smectics: effects of disorder

Stefan Scheidl and Felix von Oppen

Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Str. 77, D-50937 Köln, Germany
(February 1, 2008)

We study the effect of disorder on quantum Hall smectics within the framework of an elastic theory. Based on a renormalization group calculation, we derive detailed results for the degrees of translational and orientational order of the stripe pattern at zero temperature and carefully map out the disorder and length-scale regimes in which the system effectively exhibits smectic, nematic, or isotropic behavior. We show that disorder always leads to a finite density of free dislocations and estimate the scale on which they begin to appear.

PACS numbers: 73.40.Hm, 73.20.Mf, 46.65.+g

Recent experiments [1] discovered a pronounced anisotropy in the resistivity of the quantum Hall system near half filling of higher Landau levels (Landau level filling factor $\nu \geq 3$), suggesting that the ground state at these filling factors breaks orientational symmetry. It is believed that this is a manifestation of the formation of striped quantum Hall states whose existence had previously been predicted based on Hartree-Fock (HF) calculations [2] (for a recent review, see [3]). The existence of striped states in higher Landau levels has also been supported by exact numerical diagonalization studies [4] and by the experimental confirmation [1] of some explicit predictions [5,6] for their transport properties.

The HF calculations predict that the ground state near half filling of higher Landau levels is a unidirectional charge density wave (CDW) whose period a is of the order of the cyclotron radius [7]. In addition to orientational order, this state also breaks translational symmetry in one direction. For this reason, it is often referred to as a quantum Hall smectic state. Going beyond mean-field level by including quantum and thermal fluctuations, one expects a variety of phases with different degrees of translational and orientational order [8,5].

Here we study the effect of disorder on quantum Hall smectics at zero temperature within the framework of an elastic theory. As a function of disorder strength, we map out in detail various regimes in which the system effectively exhibits smectic (with both translational and orientational order), nematic (with orientational order only), or isotropic behavior. A concise summary of our results is provided in Fig. 1.

At low energies, the relevant gapless excitations are fluctuations of the displacement field $u(\mathbf{r}, \tau)$ associated with translational symmetry breaking [5,3] and defined as the phase of the charge-density modulation $\rho(\mathbf{r}, \tau) = \rho_0 \cos[k(r_{\parallel} + u(\mathbf{r}, \tau))]$. The position vector $\mathbf{r} = (r_{\parallel}, r_{\perp})$ is represented by its components parallel and perpendicular to the CDW wave vector $\mathbf{k}$ with $k = 2\pi/a$. We note that as opposed to the stripe position described by u , the stripe width is a massive mode that can be neglected.




FIG. 1. Illustration of regimes in dependence of disorder strength Δ_V and transverse length scale $L_{\perp}$ for $q_s \approx 0.01k$ as is typical for experiments. In the smectic regime $L_{\perp} \ll \xi_{\perp}^{\text{trans}}$ translational and orientational order are well pronounced, $f \ll k^{-2}$ and $g \ll 1$. In the weak-disorder case, $\Delta_V \ll \Delta_V^c$, this is followed by a quasi-smectic regime with quasi-long-range translational order. In the strong-disorder case, $\Delta_V \gg \Delta_V^c$, a nematic regime follows, where translational order is effectively short ranged. The strict validity of our analysis based on elastic theory ends at $L_{\perp} = L_{\perp}^{\text{disloc}}$, where free dislocations appear. Nevertheless, since in the absence of dislocations orientational order would be long-ranged (i.e. $\xi_{\perp}^{\text{orient}} = \infty$; except for extremely strong disorder), nematic order possibly persists even for $L_{\perp} \gg L_{\perp}^{\text{disloc}}$. dc transport measurements on macroscopic samples probe the regime in the indicated area.

The chiral nature of quantum Hall systems leads to a dynamic behavior of quantum Hall smectics which differs in important ways from that of ordinary smectic liquid crystals. The elastic action for quantum Hall smectics takes the form [5] (β denotes the inverse temperature)

$$S_{\text{el}}[u] = \frac{1}{\beta} \sum_{\omega} \sum_{\mathbf{q}} M \frac{\omega^2}{q_{\perp}^2} |u(\mathbf{q}, \omega)|^2 + \int_0^{\beta\hbar} d\tau E(\tau), \quad (1)$$

where M is a dynamic coefficient and $E(\tau)$ denotes the elastic free energy. The inverse powers of the transverse

wave vector $q_\perp$ in the dynamic term are *specific* to quantum Hall smectics. Most importantly, they imply that smectic order is stable against (weak) quantum fluctuations at zero temperature in the quantum Hall system, even in the absence of an alignment force orienting the stripes along a preferred direction [5].

The elastic free energy is of the smectic form [5]

$$E(\tau) = \frac{1}{2} \int_{\mathbf{r}} \left[B_\parallel \left(\nabla_\parallel u + \frac{1}{2} (\nabla_\perp u)^2 + \frac{C}{2} \right)^2 + K (\nabla_\perp^2 u)^2 + B_\perp (\nabla_\perp u)^2 \right]. \quad (2)$$

The constant C is determined by the requirement $\langle \nabla_\parallel u \rangle = 0$ which fixes the CDW wave length. The compression modulus $B_\parallel$ and the bending modulus K are related by $B_\parallel = 4k^2 K$ [9]. For $B_\perp = 0$, this free energy is rotation invariant due to the anharmonic terms [9]. We have included a crystal field with tilt modulus $B_\perp$ which accounts for the presence of an alignment force orienting the stripe pattern along a preferred direction. Experiments suggest that this alignment force is quite weak, $B_\perp \ll B_\parallel$. The presence of the tilt modulus leads to an important (transverse) length scale in the problem, defined by $q_s = (B_\perp/K)^{1/2}$. For $q_\perp \gg q_s$, the bending energy of the stripes dominates over the tilt energy while for $q_\perp \ll q_s$, the tilt energy dominates.

Various kinds of disorder may affect the quantum Hall stripes. A disorder potential $U(\mathbf{r})$ is included via an impurity action $S_{\text{imp},V} = \int_{\mathbf{r},\tau} U(\mathbf{r}) \rho(\mathbf{r},\tau)$. It can be rewritten as $S_{\text{imp},V} = (\rho_0/2) \int_{\mathbf{r},\tau} [V(\mathbf{r}) e^{iku(\mathbf{r},\tau)} + \text{c.c.}]$, where V is effectively locally correlated, $\langle V(\mathbf{r}) V^*(\mathbf{r}') \rangle \simeq W \delta(\mathbf{r} - \mathbf{r}')$, even for long-range-correlated impurity potentials $U(\mathbf{r})$ because of the oscillatory nature of the charge density $\rho(\mathbf{r},\tau)$. The effective disorder strength W is given by $W \simeq \langle U(-\mathbf{k}) U(\mathbf{k}) \rangle$. For quantitative estimates, we employ the impurity potential created by the ionized impurities (of areal density n_{imp}) in the spacer layer which dominates the transport properties of two-dimensional electron systems (2DES). For uncorrelated impurity positions, the disorder potential experienced by the electrons in the 2DES has $W = e^2 n_{\text{imp}} v_d^2(k)$. Here, $v_d(q) = 2\pi e^{-qd}/\epsilon(q)q$ is the screened Coulomb interaction, $\epsilon(q)$ the dielectric function, and d the distance of the impurity layer from the 2DES.

While the bare action contains only the point disorder $U(\mathbf{r})$, the action of the renormalization group (RG) generates a second type of disorder. This effective disorder corresponds to a random tilting force acting on the stripe pattern and is described by the additional term $S_{\text{imp},h} = \int_{\mathbf{r},\tau} h(\mathbf{r}) \nabla_\perp u(\mathbf{r},\tau)$ in the action with a local correlator $\langle h(\mathbf{r}) h(\mathbf{r}') \rangle = \Delta_h \delta(\mathbf{r} - \mathbf{r}')$.

We proceed in a standard fashion by averaging the free energy over the disorder potential using the replica trick which leads to the impurity-averaged replicated partition function

$$\langle Z^n \rangle = \int \prod_{\alpha} [du_{\alpha}] \exp \left\{ -\frac{1}{\hbar} \sum_{\alpha} S_{\text{el}}[u_{\alpha}] + \frac{1}{2\hbar^2} \int_{\mathbf{r},\tau,\tau'} \sum_{\alpha,\beta} \left[\Delta_h (\nabla_\perp u_{\alpha}(\mathbf{r},\tau)) (\nabla_\perp u_{\beta}(\mathbf{r},\tau')) + \Delta_V \cos[k(u_{\alpha}(\mathbf{r},\tau) - u_{\beta}(\mathbf{r},\tau'))] \right] \right\}. \quad (3)$$

Here, $\Delta_V = (1/2)\rho_0^2 W$ measures the strength of the point disorder while Δ_h denotes the strength of the tilt disorder.

Since translational order is stable against quantum fluctuations at zero temperature [5], one might attempt to treat the disorder potential perturbatively by expanding the action in Eq. (3) to quadratic order in u , setting $\Delta_h = 0$, and neglecting dislocations. In the resulting quadratic theory it is straight forward to calculate the correlation functions

$$f(\mathbf{r}) = \langle [u(\mathbf{r}') - u(\mathbf{r}' + \mathbf{r})]^2 \rangle, \quad (4a)$$

$$g(\mathbf{r}) = \langle [\nabla'_\perp u(\mathbf{r}') - \nabla'_\perp u(\mathbf{r}' + \mathbf{r})]^2 \rangle, \quad (4b)$$

which describe the degrees of translational and orientational order, respectively. Neglecting quantum fluctuations which are additive in perturbation theory, one finds at zero temperature $f(\mathbf{r}_\parallel) = \frac{1}{2} \int_{q_\perp} (k^2 \Delta_V / B_\parallel^2 Q_\parallel^3) [1 - (1 + Q_\parallel r_\parallel) e^{-Q_\parallel r_\parallel}]$ and $f(\mathbf{r}_\perp) = \frac{1}{2} \int_{q_\perp} (k^2 \Delta_V / B_\parallel^2 Q_\parallel^3) [1 - \cos(q_\perp r_\perp)]$. Here $\mathbf{r}_\parallel$ ($\mathbf{r}_\perp$) are vectors in the parallel (perpendicular) direction and $Q_\parallel(q_\perp) = [(B_\perp q_\perp^2 + K q_\perp^4) / B_\parallel]^{1/2}$. The analogous expressions for $g(\mathbf{r})$ differ from those for $f(\mathbf{r})$ merely by an additional factor $q_\perp^2$ in the integrand.

One readily finds that $f(\mathbf{r})$ is infrared divergent for any nonzero $\mathbf{r}$, irrespectively of whether the tilt modulus $B_\perp$ vanishes or not. Because of the multistability of the disorder potential, perturbation theory breaks down on scales where $f \gtrsim a^2$ [10], i.e. *on all finite scales*. Below, we perform a RG calculation which properly accounts for the multistability by a significant downward renormalization of the disorder strength on large scales.

The perturbative expression can nevertheless be used to estimate the characteristic transverse scale q_V , below which the renormalization sets in. To do so, we demand $f(q_V^{-1} \mathbf{e}_\perp) \sim a^2$ with the provision that the integral over $q_\perp$ in the perturbative expression for f be self-consistently restricted to $q_\perp \geq q_V$. The explicit expression for q_V depends on the disorder strength. In the *weak-disorder regime* $q_V \ll q_s$, one finds $q_V = [\Delta_V / \Delta_V^c]^{1/2} q_s$, while in the *strong-disorder regime* $q_V \gg q_s$ one obtains $q_V = [\Delta_V / \Delta_V^c]^{1/5} q_s$. The disorder strength separating these two limits is given by $\Delta_V^c = 4\pi B_\parallel^{1/2} B_\perp^{5/2} / k^4 K$.

We now turn to the RG calculation, restricting ourselves to zero temperature where the smectic state is stable in the pure system. As the approach neglects topological defects such as dislocations, the results are valid

only up to the scale $L_{\perp}^{\text{disloc}}$ where dislocations start to appear. This length will be estimated below. Our analysis builds on previous RG studies for classical smectics [9,11], adapting them to two dimensions where disorder is known to play a peculiar role [12]. We employ a conventional coarse-graining procedure, where all displacement modes in the range $\Lambda < |q_{\perp}| < k$ with $\Lambda = ke^{-l}$ are integrated out. (A length rescaling is omitted.) Treating both the point disorder and the anharmonic terms in the elastic action to one-loop order leads to the flow equations

$$\frac{d\Delta_V}{dl} = -a_1\Delta_V, \quad (5a)$$

$$\frac{d\Delta_h}{dl} = a_2\Delta_V + a_3\Delta_h, \quad (5b)$$

$$\frac{dB_{\parallel}}{dl} = -3a_3B_{\parallel}, \quad (5c)$$

$$\frac{dK}{dl} = 2a_3K, \quad (5d)$$

where $a_1(\Lambda) = k^4\Lambda\Delta_V/4\pi B_{\parallel}^2Q_{\parallel}^3(\Lambda)$ and $a_2(\Lambda) = c(k^2/\Lambda^2)a_1(\Lambda)$ with c a constant of order unity. Moreover, $a_3(\Lambda) = B_{\parallel}^{1/2}\Delta_h/16\pi K^{5/2}\Lambda^3$ for $\Lambda \gg q_s$ and $a_3 = 0$ for $\Lambda \ll q_s$. The couplings a_1 and a_2 arise from the point disorder while a_3 is due to the anharmonic terms in the elastic action. We note that the tilt modulus $B_{\perp}$ remains unrenormalized. The renormalization of $B_{\parallel}$ and K is due to the combined effect of disorder and anharmonic elasticity. The characteristic disorder scale q_V can be recovered from the flow equations by the condition $a_1(q_V) = 1$ which defines the effective onset of the renormalization of Δ_V . We first solve the flow equations ignoring the anharmonic terms in the elastic action (i.e. $a_3 = 0$). As discussed below, these terms should lead only to minor modifications. Solving the flow equation we find that the effective disorder strength $\Delta_V \sim \Lambda^2$ flows to zero for $\Lambda \ll q_V$, while the tilt disorder strength Δ_h increases logarithmically. Due to this decay of Δ_V the coupling a_1 remains finite for vanishing Λ , which indicates that the renormalization of disorder is captured consistently on one-loop level.

The detailed results depend on the disorder strength. In the *weak-disorder regime* $q_V \ll q_s$ or $\Delta_V \ll \Delta_V^c$, we find for the translational order in the soft (perpendicular) direction

$$f(\mathbf{r}_{\perp}) \sim \frac{1}{k^2} \begin{cases} (q_V r_{\perp})^2 \ln(q_s/q_V), & r_{\perp} \ll q_s^{-1}, \\ (q_V r_{\perp})^2 \ln(1/q_V r_{\perp}), & q_s^{-1} \ll r_{\perp} \ll q_V^{-1}, \\ \ln^2(q_V r_{\perp}), & r_{\perp} \gg q_V^{-1}. \end{cases} \quad (6)$$

As opposed to the perturbative result, f is now finite for any finite $r_{\perp}$ and we can identify a correlation length $\xi_{\perp}^{\text{trans}}$ for the translational order by $f(\xi_{\perp}^{\text{trans}}) = a^2$ and find $\xi_{\perp}^{\text{trans}} \sim q_V^{-1}$. The slow $\ln^2$ increase of f in the regime

$r_{\perp} \gg \xi_{\perp}^{\text{trans}}$, typical of two-dimensional disordered systems [12], implies that the system exhibits “quasi”-long-range translational order, with power-law Bragg peaks with logarithmically varying powers. The orientational correlation function g remains finite, $g(\mathbf{r}_{\perp}) \sim (q_V/k)^2 \ln(q_s/q_V) \ll 1$, for arbitrarily large lengths $r_{\perp} \gg q_V^{-1}$. Hence the correlation length $\xi_{\perp}^{\text{orient}}$ for orientational order, defined by $g(\xi_{\perp}^{\text{orient}}) = 1$, is infinitely large.

For sufficiently strong disorder, we enter the *strong-disorder regime* $q_V \gg q_s$ or $\Delta_V \gg \Delta_V^c$ where

$$f(\mathbf{r}_{\perp}) \sim \frac{1}{k^2} \begin{cases} (q_V/q_s)^3 (q_s r_{\perp})^2, & r_{\perp} \ll q_s^{-1}, \\ (q_V/q_s)^3 \ln(q_s r_{\perp}), & q_s^{-1} \ll r_{\perp} \ll q_h^{-1}, \\ \ln^2(q_s r_{\perp}), & r_{\perp} \gg q_h^{-1}. \end{cases} \quad (7)$$

In this case, the $\ln^2$ regime typically sets in only at extremely large length scales $r_{\perp} \gg q_h^{-1}$ with $q_h = q_s \exp[-(q_V/q_s)^3]$ so that we can restrict attention to the $\ln$ regime. At first sight, this seems to suggest the presence of quasi-long-range translational order in this case. While this is formally true, the corresponding singularities in the structure factor are extremely weak due to the large prefactor $(q_V/q_s)^3$ and for practical purposes, there is only short-range translational order with correlation length $\xi_{\perp}^{\text{trans}} = (q_s/q_V)^{1/2} q_V^{-1}$. As expected, the degree of translational order is significantly reduced compared to the weak-disorder regime. The degree of orientational order is also reduced, since $g(\mathbf{r}_{\perp}) \sim (q_V/k)^2 (q_V/q_s)$ for $r_{\perp} \gg q_s^{-1}$. Nevertheless, $\xi_{\perp}^{\text{orient}} = \infty$ as long as $q_V^3 \ll k^2 q_s$. Otherwise, for extremely strong disorder, $\xi_{\perp}^{\text{orient}} \sim k^2/q_V^3$.

Results for the correlation functions in the parallel direction in the asymptotic regimes can be easily obtained from those in the perpendicular direction by the replacement $r_{\perp} \rightarrow (q_s/k)r_{\parallel}$. The small factor $(q_s/k) \ll 1$ reflects the fact that the system is more rigid in the parallel than in the perpendicular direction. It is interesting to note that in the absence of an alignment force ($B_{\perp} = 0$) orienting the stripes along a preferred direction, disorder destroys translational order and leads to quasi-long-range orientational order, even when neglecting dislocations.

So far the density modulation has been treated assuming the absence of any topological defects. We now show that disorder always induces a finite density of dislocations and estimate the transverse scale $L_{\perp}^{\text{disloc}}$ on which dislocations become relevant and which limits the range of validity of the above results. The occurrence of dislocations depends on a competition between the elastic energy cost E_{self} to create a dislocation and the energy gain E_{pin} in the disorder potential. The energy gain E_{pin} originates from the reduced rigidity of the charge density profile in the presence of dislocations. The scale $L_{\perp}^{\text{disloc}}$ is identified as the size of a finite system where the free energy $E_{\text{self}} + E_{\text{pin}}$ of a single dislocation becomes negative. For a strictly smectic system ($B_{\perp} = 0$), the elastic energy cost E_{self} of a dislocation is known to be finite.

A finite tilt modulus $B_\perp$ leads to a logarithmic increase in the elastic energy with system size $L_\perp$ for $L_\perp \gg q_s^{-1}$. The energy of the dislocation in the disorder potential is a random variable that depends on the location of the dislocation. The pinning energy E_{pin} is given by the *maximal* energy that the dislocation can gain from the disorder potential. Assuming a Gaussian distribution for the disorder energy, this energy can be computed using the statistics of extreme values. In the *weak-disorder regime*, we eventually find that the creation of dislocations becomes favorable on transverse scales

$$L_\perp^{\text{disloc}} \sim q_V^{-1} \exp[c \ln^{2/3}(q_s/q_V)] \quad (8)$$

(with c a constant of order unity) corresponding to an areal dislocation density of $n_{\text{disloc}} \sim (L_\perp^{\text{disloc}})^{-2} (B_\perp/B_\parallel)^{1/2}$. In the *strong-disorder limit*, we find

$$L_\perp^{\text{disloc}} \sim q_s^{-1} \min[1, k^2 q_s/q_V^3] \quad (9)$$

and a dislocation density of $n_{\text{disloc}} \sim (L_\perp^{\text{disloc}})^{-3} (K/B_\parallel)^{1/2}$. Thus, within a finite range of disorder strengths, $\Delta_V^c \ll \Delta_V \ll (k/q_s)^{10/3} \Delta_V^c$ or $q_s^3 \ll q_V^3 \ll k^2 q_s$, $L_\perp^{\text{disloc}}$ saturates at q_s^{-1} , before it shrinks further, coinciding with $\xi_\perp^{\text{orient}}$.

The present analysis of regimes neglects the influence of the anharmonic elastic terms, although they are included at one-loop level in the flow equations (5). On scales $q_\perp \ll q_V$ these terms lead to an effective reduction of $B_\parallel$ and an increase of K , making the system appear more isotropic. The integration of the full flow equations [13] modifies the correlation functions *only slightly* in the strong-disorder limit. In addition, q_s is reduced due to the renormalization of K .

Our results therefore lead to the following overall picture, as summarized in Fig. 1. In the clean limit we have a sequence of transverse length scales $q_s^{-1} \ll q_V^{-1} \simeq \xi_\perp^{\text{trans}} \ll L_\perp^{\text{disloc}}$ (see Fig. 1). On scales $L_\perp \ll q_V^{-1}$, the system looks like a *smectic* with pronounced translational order ($f \ll a^2$) and no free dislocations. On scales $q_V^{-1} \ll L_\perp \ll L_\perp^{\text{disloc}}$, there are still no free dislocations. However, $f \gg a^2$ and f increases logarithmically, displaying quasi-long-range order, whence we call this regime *quasi-smectic*. For $L_\perp \gg L_\perp^{\text{disloc}}$ dislocations appear and destroy translational order. Neglecting dislocations, one finds $g \ll 1$ (i.e. $\xi_\perp^{\text{orient}} = \infty$) on all scales in the weak-disorder regime, which indicates the possibility to have good orientational (nematic) order even for $L_\perp \gg L_\perp^{\text{disloc}}$.

In the strong-disorder regime, the sequence of scales is $\xi_\perp^{\text{trans}} \ll q_V^{-1} \ll L_\perp^{\text{disloc}} \lesssim q_s$. Good translational order survives only for $L_\perp \ll \xi_\perp^{\text{trans}}$. Beyond this scale, translational order is rapidly lost as f increases as a power law with $f \gg 1$. On the other hand, orientational order is still retained, $g \ll 1$. Hence, we cross over from a *smectic* regime for $L_\perp \ll \xi_\perp^{\text{trans}}$ to a *nematic* regime for $L_\perp \gg \xi_\perp^{\text{trans}}$. Dislocations are present

on scales $L_\perp \gg L_\perp^{\text{disloc}}$. For $\Delta_V^c \ll \Delta_V \ll (k/q_s)^{10/3} \Delta_V^c$, $\xi_\perp^{\text{orient}} = \infty$ in the absence of dislocations and nematic order may survive beyond $L_\perp^{\text{disloc}}$ even in their presence. For $\Delta_V \gg (k/q_s)^{10/3} \Delta_V^c$, $\xi_\perp^{\text{orient}} \sim L_\perp^{\text{disloc}}$ and most likely nematic order is destroyed for $L_\perp \gg \xi_\perp^{\text{orient}}$, leading to an isotropic regime.

For typical experimental samples [1], we estimate $q_s \sim q_V \sim 0.01k$, indicating that they are located in the crossover region between the weak- and strong-disorder regimes (cf. Fig. 1). While we cannot firmly deduce the presence of orientational order on the macroscopic scales probed by *dc* transport measurements, there is a good chance that orientational order persists even for $L_\perp \gtrsim L_\perp^{\text{disloc}}$, since $\xi_\perp^{\text{orient}} = \infty$ for $\Delta_V \ll (k/q_s)^{10/3} \Delta_V^c$ in the absence of dislocations. The regimes identified in this paper at smaller lengths may be accessible to other experimental probes such as surface acoustic waves.

This work was partly supported by the DFG through Schwerpunktprogramm ‘‘Quanten-Hall-Systeme’’ and Sonderforschungsbereich 341.

-
- [1] M. P. Lilly *et al.*, Phys. Rev. Lett. **82**, 394 (1999); R. Du *et al.*, Solid State Comm. **109**, 389 (1999); J.P. Eisenstein *et al.*, cond-mat/0003405.
 - [2] M. M. Fogler, A. A. Koulakov, and B. I. Shklovskii, Phys. Rev. B **54**, 1853 (1996); R. Moessner and J. T. Chalker, Phys. Rev. B **54**, 5006 (1996).
 - [3] F. von Oppen, B. I. Halperin, and A. Stern, cond-mat/0002087.
 - [4] E. H. Rezayi, F. D. M. Haldane, and K. Yang, Phys. Rev. Lett. **83**, 1219 (1999).
 - [5] A.H. MacDonald and M. P. A. Fisher, Phys. Rev. B **61**, 5724 (2000).
 - [6] F. von Oppen, B.I. Halperin, and A. Stern, Phys. Rev. Lett. **84**, 2937 (2000).
 - [7] In principle, the CDW is unstable with respect to charge ordering along the stripes at extremely low temperatures [5]. Presumably this ordering is destroyed by minute impurity concentrations even at zero temperature. Thus we will ignore this complication.
 - [8] E. Fradkin and S. A. Kivelson, Phys. Rev. B **59**, 8065 (1999).
 - [9] G. Grinstein, and R. A. Pelcovits, Phys. Rev. A **26**, 915 (1982).
 - [10] H. Fukuyama and P. A. Lee, Phys. Rev. B **17**, 535 (1978); P. A. Lee and T. M. Rice, Phys. Rev. B **19**, 3970, (1979); A. I. Larkin, Sov. Phys. JETP **31**, 784 (1970).
 - [11] L. Radzihovsky and J. Toner, Phys. Rev. B **60**, 206 (1999).
 - [12] J. Villain and J.F. Fernandez, Z. Phys. B **54**, 139 (1984); for a recent review, see T. Nattermann and S. Scheidl, Adv. Phys. **49**, 607 (2000).
 - [13] S. Scheidl and F. von Oppen, to be published.